

AD-A262 297



2

NPS-MA-93-009

NAVAL POSTGRADUATE SCHOOL

Monterey, California



DTIC
SELECTE
MAR 17 1993
S B D

93-05470



2384

A PARALLEL DIVIDE AND CONQUER
ALGORITHM FOR THE GENERALIZED REAL
SYMMETRIC DEFINITE TRIDIAGONAL
EIGENPROBLEM

by

Carlos F. Borges
William B. Gragg

Technical Report For Period
September 1992 - December 1992

Approved for public release; distribution unlimited

Prepared for: Naval Postgraduate School
Monterey, CA 93943

93 3 16 091

NAVAL POSTGRADUATE SCHOOL
MONTEREY, CA 93943

Rear Admiral R. W. West, Jr.
Superintendent


Harrison Shull
Provost

This report was prepared in conjunction with research conducted for the Naval Postgraduate School and funded by the Naval Postgraduate School.


Reproduction of all or part of this report is authorized.

This report was prepared by:


CARLOS BORGES
Assistant Professor


WILLIAM B. GRAGG
Professor

Reviewed by:


RICHARD FRANKE
Chairman
Department of Mathematics

Released by:


PAUL J. MARTO
Dean of Research

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NPS-MA-93-009			5. MONITORING ORGANIZATION REPORT NUMBER(S) NPS-MA-93-009		
6a. NAME OF PERFORMING ORGANIZATION Naval Postgraduate School		6b. OFFICE SYMBOL (if applicable) MA	7a. NAME OF MONITORING ORGANIZATION Naval Postgraduate School		
6c. ADDRESS (City, State, and ZIP Code) Monterey, CA 93943			7b. ADDRESS (City, State, and ZIP Code) Monterey, CA 93943		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Naval Postgraduate School		8b. OFFICE SYMBOL (if applicable) MA	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER O&MN Direct Funding		
8c. ADDRESS (City, State, and ZIP Code) Monterey, CA 93943			10. SOURCE OF FUNDING NUMBERS		
		PROGRAM ELEMENT NO	PROJECT NO	TASK NO	WORK UNIT ACCESSION NO
11. TITLE (Include Security Classification) A Parallel Divide Conquer Algorithm for the Generalized Real Symmetric Definite Tridiagonal Eigenproblem					
12. PERSONAL AUTHOR(S) Carlos F. Borges, William B. Gragg					
13a. TYPE OF REPORT Technical Report		13b. TIME COVERED FROM 9-92 TO 12-92		14. DATE OF REPORT (Year, Month, Day) 921216	
15. PAGE COUNT 19					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Parallel divide and conquer algorithm, Tridiagonal eigenproblem, eigenvectors		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) We develop a parallel divide and conquer algorithm, by extension, for the generalized real symmetric definite tridiagonal eigenproblem. The algorithm employs techniques first proposed by Gu and Eisenstat to prevent loss of orthogonality in the computed eigenvectors for the modification algorithm. We examine numerical stability and adapt the insightful error analysis of Gu and Eisenstat to the arrow case. The algorithm incorporates an elegant zero finder with global monotone cubic convergence that has performed well in numerical experiments. A complete set of tested matlab routines implementing the algorithm is available on request from the authors.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Carlos Borges			22b. TELEPHONE (Include Area Code) (408)656-2124		22c. OFFICE SYMBOL MA/Bc

DD Form 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

S/N 0102-LF-014-6603

A Parallel Divide and Conquer Algorithm for the Generalized Real Symmetric Definite Tridiagonal Eigenproblem

Carlos F. Borges and William B. Gragg†*

Abstract.

We develop a parallel divide and conquer algorithm, by *extension*, for the generalized real symmetric definite tridiagonal eigenproblem. The algorithm employs techniques first proposed by Gu and Eisenstat to prevent loss of orthogonality in the computed eigenvectors for the *modification* algorithm. We examine numerical stability and adapt the insightful error analysis of Gu and Eisenstat to the arrow case. The algorithm incorporates an elegant zero finder with global monotone cubic convergence that has performed well in numerical experiments. A complete set of tested matlab routines implementing the algorithm is available on request from the authors.

1 Introduction

We consider the problem of finding a matrix $U \in \mathbb{R}^{n \times n}$ such that

$$U^T (T - S\lambda) U \equiv \Lambda - I\lambda,$$

is diagonal, or equivalently

$$U^T S U = I \quad \text{and} \quad U^T T U = \Lambda, \tag{1}$$

where

*Authors address: Code Ma/Bc, Naval Postgraduate School, Monterey, CA 93943. Email: borges@waylon.math.nps.navy.mil

†Authors address: Code Ma/Gr, Naval Postgraduate School, Monterey, CA 93943. Email: gragg@guinness.math.nps.navy.mil

$$T = \begin{bmatrix} \alpha_1 & \beta_1 & & & \\ \beta_1 & \alpha_2 & \beta_2 & & \\ & \beta_2 & & \ddots & \\ & & \ddots & \beta_{n-1} & \\ & & & \beta_{n-1} & \alpha_n \end{bmatrix}, \quad S = \begin{bmatrix} \delta_1 & \gamma_1 & & & \\ \gamma_1 & \delta_2 & \gamma_2 & & \\ & \gamma_2 & & \ddots & \\ & & \ddots & \gamma_{n-1} & \\ & & & \gamma_{n-1} & \delta_n \end{bmatrix},$$

and S is assumed to be positive definite. This generalized eigenvalue problem has two special cases that are of interest in themselves. They are:

1. $S = I$, the ordinary tridiagonal eigenproblem.
2. $S = I$ and $\alpha_k \equiv 0$, the bidiagonal singular value problem (BSVP), by *perfect shuffle* of the Jordan matrix

$$\begin{bmatrix} 0 & B^T \\ B & 0 \end{bmatrix}$$

with B upper bidiagonal [16].

There are two phases to the divide and conquer algorithm, the divide (or split) phase, and the conquer (or consolidate) phase. We shall address these in order.

2 The algorithm

2.1 The divide phase

Denote by e_i the i th axis vector where the dimension will be clear from the context. Let $s, 1 \leq s \leq n$, be an integer, the *split index*, and consider the following block forms:

$$T = \begin{bmatrix} T_1 & e_{s-1}\beta_{s-1} & \\ \beta_{s-1}e_{s-1}^T & \alpha_s & \beta_s e_1^T \\ & e_1\beta_s & T_2 \end{bmatrix},$$

$$S = \begin{bmatrix} S_1 & e_{s-1}\gamma_{s-1} & \\ \gamma_{s-1}e_{s-1}^T & \delta_s & \gamma_s e_1^T \\ & e_1\gamma_s & S_2 \end{bmatrix}.$$

Note that $s = n$ is possible; then T_2 , S_2 , and e_1 are empty [9, 10]. Suppose we solve the subproblems

$$U_k^T (T_k - S_k \lambda) U_k \equiv \Lambda_k - I \lambda \quad (k = 1, 2). \quad (2)$$

The form of the subproblems is preserved. In particular, the matrices S_k are positive definite and, if T has a zero diagonal, so do the matrices T_k . Let

$$\hat{U} = \begin{bmatrix} U_1 & & \\ & 1 & \\ & & U_2 \end{bmatrix}.$$

Then

$$\hat{U}^T (T - S\lambda) \hat{U} = \begin{bmatrix} U_1^T (T_1 - S_1 \lambda) U_1 & U_1^T e_{s-1} (\beta_{s-1} - \gamma_{s-1} \lambda) & \\ (\beta_{s-1} - \gamma_{s-1} \lambda) e_{s-1}^T U_1 & \alpha_s - \delta_s \lambda & (\beta_s - \gamma_s \lambda) e_1^T U_2 \\ & U_2^T e_1 (\beta_s - \gamma_s \lambda) & U_2^T (T_2 - S_2 \lambda) U_2 \end{bmatrix}.$$

2.2 The conquer phase

The conquer phase consists of solving the subproblems (2) from the divide phase, consolidating the solutions, and finally, solving the consolidated problem. Let

$$u_1 = U_1^T e_{s-1}, \quad u_2 = U_2^T e_1,$$

where the U_k are solutions to (2). Then

$$\hat{U}^T (T - S\lambda) \hat{U} = \begin{bmatrix} \Lambda_1 - I\lambda & u_1(\beta_{s-1} - \gamma_{s-1}\lambda) & \\ (\beta_{s-1} - \gamma_{s-1}\lambda)u_1^T & \alpha_s - \delta_s\lambda & (\beta_s - \gamma_s\lambda)u_2^T \\ & u_2(\beta_s - \gamma_s\lambda) & \Lambda_2 - I\lambda \end{bmatrix}.$$

The right side is the sum of a diagonal and a *Swiss cross*:

$$\hat{U}^T (T - S\lambda) \hat{U} = \begin{bmatrix} x & & & & \\ & x & & & \\ & & x & & \\ & & & x & \\ & & & & x \end{bmatrix} + \begin{bmatrix} & & x & & \\ & & x & & \\ x & x & x & x & x \\ & & x & & \\ & & x & & \end{bmatrix}.$$

This can be permuted to an arrow matrix by a permutation similarity transformation with $P_s = [e_1, e_2, \dots, e_{s-1}, e_{s+1}, \dots, e_n, e_s]$. Thus

$$\begin{aligned} \hat{A}(\lambda) &:= P_s^T \hat{U}^T (T - S\lambda) \hat{U} P_s \\ &= \begin{bmatrix} \Lambda_1 & & u_1 \beta_{s-1} \\ & \Lambda_2 & u_2 \beta_s \\ \beta_{s-1} u_1^T & \beta_s u_2^T & \alpha_s \end{bmatrix} - \begin{bmatrix} I & & u_1 \gamma_{s-1} \\ & I & u_2 \gamma_s \\ \gamma_{s-1} u_1^T & \gamma_s u_2^T & \delta_s \end{bmatrix} \lambda \\ &= \begin{bmatrix} D & Bu \\ u^T B & \alpha \end{bmatrix} - \begin{bmatrix} I & Cu \\ u^T C & \gamma \end{bmatrix} \lambda, \end{aligned}$$

with

Session For	
US GRA&I	<input checked="" type="checkbox"/>
IC TAB	<input type="checkbox"/>
unannounced	<input type="checkbox"/>
Classification	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

DTIC QUALITY INSPECTED 1/

$$u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix},$$

$$B = \begin{bmatrix} \beta_{s-1}I & \\ & \beta_s I \end{bmatrix}, \quad C = \begin{bmatrix} \gamma_{s-1}I & \\ & \gamma_s I \end{bmatrix}.$$

Since S and $\begin{bmatrix} I & Cu \\ u^T C & \gamma \end{bmatrix}$ are congruent the latter inherits positive definiteness from the former. Its Cholesky decomposition is

$$\begin{bmatrix} I & Cu \\ u^T C & \gamma \end{bmatrix} = \begin{bmatrix} I & \\ u^T C & \rho \end{bmatrix} \begin{bmatrix} I & Cu \\ & \rho \end{bmatrix} = R^T R,$$

with $\rho^2 = \gamma - u^T C^2 u > 0$ the Schur complement in S of

$$\begin{bmatrix} S_1 & \\ & S_2 \end{bmatrix}.$$

Now

$$R^{-1} = \begin{bmatrix} I & -Cu/\rho \\ & 1/\rho \end{bmatrix}$$

and a second congruence transformation with R^{-1} gives

$$\begin{aligned} A(\lambda) &:= R^{-T} \hat{A}(\lambda) R^{-1} \\ &= R^{-T} \begin{bmatrix} D & Bu \\ u^T B & \alpha \end{bmatrix} R^{-1} - I\lambda \\ &= \begin{bmatrix} D & w \\ w^T & \omega \end{bmatrix} - I\lambda =: A - \lambda I \end{aligned}$$

with

$$\begin{aligned} w &= \frac{(B - DC)u}{\rho}, \\ \omega &= \frac{\alpha_s - u^T (2B - DC)Cu}{\rho^2}. \end{aligned}$$

We have reduced the conquer step to the problem of solving an ordinary eigenproblem for a symmetric arrow matrix. If V is an orthogonal matrix with

$$AV = V\Lambda$$

and Λ diagonal, then (1) holds with

$$\begin{aligned} U &= \hat{U} P_s R^{-1} V \\ &= \begin{bmatrix} U_1 & -U_1 u_1 \gamma_{s-1}/\rho \\ & 1/\rho \\ & U_2 & -U_2 u_2 \gamma_s/\rho \end{bmatrix} V. \end{aligned}$$

It is useful that $\mathbf{v}_k = U_k \mathbf{u}_k$ can be computed in $O(n)$ time by solving $S_1 \mathbf{v}_1 = \mathbf{e}_{s-1}$ and $S_2 \mathbf{v}_2 = \mathbf{e}_1$ using the Cholesky factorization $S_1 = L_1 L_1^T$ and the reverse Cholesky factorization $S_2 = L_2^T L_2$. In the case that only the eigenvalues are wanted it is only necessary to compute the first and last rows of the U -matrices which constitutes a further savings.

In summary, the conquer phase proceeds by consolidating the subproblems and building a full eigenproblem for an arrow matrix.

3 Solving the eigenproblem for the arrow

In this section we consider the solution of the eigenproblem for a real symmetric arrow matrix

$$A = \begin{bmatrix} D & \mathbf{b} \\ \mathbf{b}^T & \gamma \end{bmatrix}$$

where $A \in \mathbb{R}^{n \times n}$ is symmetric, $D = \text{diag}(\mathbf{a})$, $\mathbf{a} = [\alpha_1, \dots, \alpha_{n-1}]^T$, $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_{n-1}$, and $\mathbf{b} = [\beta_1, \dots, \beta_{n-1}]^T \geq 0$. When A arises from the BSVD then \mathbf{a} is odd and \mathbf{b} is even, that is $\mathbf{a} + J\mathbf{a} = 0$ and $\mathbf{b} = J\mathbf{b}$, with J the *counter-identity*, the identity matrix with its columns reversed, and $\gamma = 0$.

If any $\beta_j = 0$ then it is possible to set $\lambda_j = \alpha_j$ and deflate the matrix since \mathbf{e}_j is clearly an eigenvector [28]. We shall call this β -deflation and note that if $\beta_j < \text{tol}_\beta \|\mathbf{b}\|$ where tol_β is a small tolerance then a *numerical* deflation occurs. We derive a precise value for tol_β in section 4.4.

A second type of deflation occurs if applying a 2×2 rotation similarity transformation in the $(j, j+1)$ -plane that takes β_j to zero introduces a sufficiently small element in the $(j, j+1)$ position of the matrix. This will be called a *comba*-deflation (see [15]). At each consolidation step we perform a sweep to check for β -deflations followed by a sweep to check for *comba*-deflations. The *comba*-deflation can be arranged so that the ordering of the α_j is preserved whenever one occurs. After deflation the new $\beta_{j+1}^* := \sqrt{\beta_j^2 + \beta_{j+1}^2} \geq \beta_{j+1}$ and hence no further β -deflation can occur. The *comba*-deflations can be disposed of with a single pass by backing up a single element whenever one occurs. Note that deflation is backward stable since it results in *small* backward errors in A . Deflation for the BSVD is more delicate involving a simultaneous sweep from both ends of the matrix. Care must be exercised at the center of the matrix.

After deflation the resulting matrix can be taken to have all $\beta_j > 0$ and the elements of the arrow *shaft* distinct and ordered, that is $\alpha_1 > \alpha_2 > \dots > \alpha_{n-1}$. An arrow matrix of this form will be called *ordered* and *reduced*. Henceforth, we shall assume A is of this form.

The block Gauss factorization of $A - \lambda I$ is

$$A - \lambda I = \begin{bmatrix} I & 0 \\ \mathbf{b}^T(D - \lambda I)^{-1} & 1 \end{bmatrix} \begin{bmatrix} D - \lambda I & \mathbf{b} \\ 0^T & -f(\lambda) \end{bmatrix}$$

where f , the *spectral function* of A , is given by

$$f(\lambda) = \lambda - \gamma + \sum_{j=1}^{n-1} \frac{\beta_j^2}{\alpha_j - \lambda}.$$

This is a rational Pick function with a pole at infinity [1]. The *most general form* of a rational Pick function is

$$f(\lambda) = \delta\lambda - \gamma + \sum_{j=1}^{n-1} \frac{\beta_j^2}{\alpha_j - \lambda}, \quad \delta \geq 0. \quad (3)$$

In relation to the various divide and conquer schemes, the case $\delta > 0$ corresponds with *extension*, $\delta = 0$ with *modification*, and $\delta = \gamma = 0$ with *restriction* [7].

Inspection of the graph of the spectral function reveals that the elements of the shaft interlace the eigenvalues

$$\lambda_1 > \alpha_1 > \lambda_2 > \dots > \alpha_{n-1} > \lambda_n. \quad (4)$$

Moreover, in the present case, the derivative of the spectral function is bounded below by one so that its zeros are, in a certain sense, well determined.

3.1 The zero finder

The fundamental problem in finding the eigenvalues of an arrow is that of providing a stable and efficient method for finding the zeros of the spectral function. We now examine this problem in some detail.

The zero finding algorithm we present is globally convergent in the sense that the iteration will converge to the unique zero of f in (α_k, α_{k-1}) from any starting value in the closed interval $[\alpha_k, \alpha_{k-1}]$, where we put $\alpha_0 = +\infty$ and $\alpha_n = -\infty$. The zero finder converges monotonically at a *cubic* rate and applies to a general Pick function as given in formula (3).

3.2 Interior intervals

The iterative procedure for finding the unique zero of f in one of the interior intervals (α_k, α_{k-1}) proceeds as follows. Let x_0 , $\alpha_k < x_0 < \alpha_{k-1}$, be an initial approximation to λ_k . If x_j is known choose

$$\phi_j(x) = \sigma + \frac{\omega_0}{\alpha_{k-1} - x} + \frac{\omega_1}{\alpha_k - x}$$

so that

$$\phi_j^{(i)}(x_j) = f^{(i)}(x_j), \quad i = 0, 1, 2. \quad (5)$$

Thus σ , ω_0 , and ω_1 must satisfy

$$\begin{bmatrix} 1 & (\alpha_{k-1} - x_j)^{-1} & (\alpha_k - x_j)^{-1} \\ 0 & (\alpha_{k-1} - x_j)^{-2} & (\alpha_k - x_j)^{-2} \\ 0 & (\alpha_{k-1} - x_j)^{-3} & (\alpha_k - x_j)^{-3} \end{bmatrix} \begin{bmatrix} \sigma \\ \omega_0 \\ \omega_1 \end{bmatrix} = \begin{bmatrix} f(x_j) \\ f'(x_j) \\ f''(x_j) \end{bmatrix}$$

and we find

$$\begin{aligned} \sigma &= 3x_j - (\gamma + \alpha_{k-1} + \alpha_k) + \sum_{i \neq k-1, k} \frac{\beta_i^2}{\alpha_i - x_j} \frac{\alpha_i - \alpha_{k-1}}{\alpha_i - x_j} \frac{\alpha_i - \alpha_k}{\alpha_i - x_j}, \\ \omega_0 &= \beta_{k-1}^2 + \frac{(\alpha_{k-1} - x_j)^3}{\alpha_{k-1} - \alpha_k} \left(1 + \sum_{i \neq k-1, k} \frac{\beta_i^2}{(\alpha_i - x_j)^2} \frac{\alpha_i - \alpha_k}{\alpha_i - x_j} \right), \\ \omega_1 &= \beta_k^2 + \frac{(x_j - \alpha_k)^3}{\alpha_{k-1} - \alpha_k} \left(1 + \sum_{i \neq k-1, k} \frac{\beta_i^2}{(\alpha_i - x_j)^2} \frac{\alpha_i - \alpha_{k-1}}{\alpha_i - x_j} \right). \end{aligned}$$

Since $\omega_0 > 0$ and $\omega_1 > 0$ it follows that ϕ_j is a Pick function. Thus ϕ_j has a unique zero $x_{j+1} \in (\alpha_k, \alpha_{k-1})$. Also

$$\omega_0 > \beta_{k-1}^2 > 0, \quad \omega_1 > \beta_k^2 > 0.$$

The error function

$$f(x) - \phi(x) = x - (\gamma + \sigma) + \sum_{i \neq k-1, k} \frac{\beta_i^2}{\alpha_i - x} + \frac{\beta_{k-1}^2 - \omega_0}{\alpha_{k-1} - x} + \frac{\beta_k^2 - \omega_1}{\alpha_k - x}$$

has n zeros, counting multiplicities. There are $n-3$ zeros exterior to (α_k, α_{k-1}) and three more at x_j . Thus the error function crosses zero exactly once in the interval (α_k, α_{k-1}) . Hence x_{j+1} lies between x_j and λ_k , and the iteration is monotonically convergent from any starting guess $x_0 \in [\alpha_k, \alpha_{k-1}]$ as claimed. The cubic rate of convergence follows from (5).

Successive iterates can be found by solving quadratic equations. Rather than solve $\phi_j(x) = 0$ for x_{j+1} it is better to solve

$$\phi_j(x_j - \Delta) = 0$$

for the increment $\Delta = x_j - x_{j+1}$. Some rearrangement using (5) reduces this to

$$\alpha \Delta^2 + \beta \Delta - f = 0, \quad (6)$$

with

$$\alpha = \frac{\sigma}{(\alpha_{k-1} - x_j)(x_j - \alpha_k)}, \quad (7)$$

$$\beta = f'(x_j) - \left(\frac{1}{\alpha_{k-1} - x_j} + \frac{1}{\alpha_k - x_j} \right) f(x_j). \quad (8)$$

When shifts of the origin to the nearest pole [15] are used then one of α_{k-1} or α_k is zero. The computation of $\beta = \beta(x_j)$ should account for the fact that it has only simple poles at α_{k-1} and α_k .

If we start at the midpoint of the interval, $x_0 = (\alpha_{k-1} + \alpha_k)/2$, then we always have $\beta = \beta(x_j) \geq f'(x_j) \geq 1$. This can be seen by noting that $\beta(x_0) = f'(x_0)$ and that when $x_0 > \lambda_k$ then for all of the succeeding iterates $f(x_j) > 0$, by monotonicity, and $\frac{1}{\alpha_{k-1} - x_j} + \frac{1}{\alpha_k - x_j}$ is negative. If $x_0 < \lambda_k$ a similar argument applies. It follows that the increment can *always* be computed stably as

$$\Delta = \frac{2f/\beta}{1 + \sqrt{1 + \frac{2\alpha}{\beta} \frac{2f}{\beta}}}. \quad (9)$$

3.3 Exterior intervals

The treatment of the two exterior intervals is *geometrically* the same as above. Again, the approximating function has poles at the endpoints and the residues at these poles, and the constant term, are chosen to satisfy (5). We present the case for the interval (α_1, ∞) , the case for the other exterior interval being similar. Now

$$\phi_j(x) = \omega_0 x - \sigma + \frac{\omega_1}{\alpha_1 - x}$$

with

$$\begin{aligned} \omega_0 &= 1 + \sum_{i=2}^{n-1} \frac{\beta_i^2}{(x_j - \alpha_i)^2} \frac{\alpha_1 - \alpha_i}{x_j - \alpha_i} \geq 1, \\ \omega_1 &= \beta_1^2 + \sum_{i=2}^{n-1} \beta_i^2 \left(\frac{x_j - \alpha_1}{x - \alpha_i} \right)^3 \geq \beta_1^2. \end{aligned}$$

The inequalities are strict unless $n = 2$. Again we find (6) where now

$$\begin{aligned} \alpha &= - \frac{1 + \sum_{i=2}^{n-1} \frac{\beta_i^2}{(x_j - \alpha_i)^2} \frac{\alpha_1 - \alpha_i}{x_j - \alpha_i}}{x_j - \alpha_1}, \\ \beta &= f'(x_j) + \frac{f(x_j)}{x_j - \alpha_1}. \end{aligned}$$

These are limiting cases of (7) and (8) (introduce another pole $\alpha_0 > \alpha_1$ and let $\alpha_0 \rightarrow +\infty$). If $x_0 > \lambda_1$ then $f(\lambda) > 0$ so $\beta > f' > 1$ and Δ is again computed stably using (9). We obtain global monotone cubic convergence as before.

Contrary to the algorithms of [11, 12, 15] our algorithm is well-defined when starting at the endpoints of the intervals. The algorithm of [23] can start at the endpoints but has only quadratic convergence.

To guarantee that $x_0 \geq \lambda_1$ we take x_0 to be the iterate in $(\alpha_1, +\infty)$ from $+\infty$. As $x_0 \rightarrow +\infty$ the approximate Pick function tends to

$$\phi(x) = x - \gamma + \frac{\|b\|^2}{\alpha_1 - x}. \quad (10)$$

Our *actual* starting guess is the zero of (10) in $(\alpha_1, +\infty)$:

$$x_0 = \begin{cases} \alpha_1 + \frac{\gamma - \alpha_1}{2} + \sqrt{\left(\frac{\gamma - \alpha_1}{2}\right)^2 + \|b\|^2} & , \quad \gamma > \alpha_1, \\ \alpha_1 + \frac{\|b\|^2}{\frac{\alpha_1 - \gamma}{2} + \sqrt{\left(\frac{\gamma - \alpha_1}{2}\right)^2 + \|b\|^2}} & , \quad \gamma \leq \alpha_1. \end{cases}$$

When shifts are used we have $\alpha_1 = 0$.

3.4 Orthogonality of the eigenvectors

It is essential that the computed eigenvectors of the arrow matrix be numerically orthogonal. As a point of entry into the further analysis of the algorithm we now examine the orthogonality of the eigenvectors following [15].

Consider the divided difference

$$\begin{aligned} f(\lambda, \mu) &= \frac{f(\lambda) - f(\mu)}{\lambda - \mu} = 1 + \sum_{j=1}^{n-1} \frac{\beta_j^2}{(\alpha_j - \lambda)(\alpha_j - \mu)} \\ &= 1 + b^T (D - \lambda I)^{-1} (D - \mu I)^{-1} b. \end{aligned} \quad (11)$$

Note that $\mu = \lambda$ gives $f'(\lambda) = 1 + \|(D - \lambda I)^{-1} b\|_2^2$. If $f(\lambda) = 0$ then

$$v(\lambda) = \begin{bmatrix} \frac{\beta_1}{\lambda - \alpha_1} \\ 1 \end{bmatrix} = \begin{bmatrix} (\lambda I - D)^{-1} b \\ 1 \end{bmatrix}$$

is an eigenvector of the arrow matrix $A = \begin{bmatrix} D & b \\ b^T & \gamma \end{bmatrix}$ associated with the eigenvalue λ , and

$$u(\lambda) = \frac{v(\lambda)}{\sqrt{f'(\lambda)}}$$

is the normalized eigenvector whose last element is positive. The ordering of A implies that its matrix of eigenvectors can be taken positive below and on the diagonal, and negative above.

Let $f(\lambda_0) = f(\mu_0) = 0$ with $\lambda_0 \neq \mu_0$. Thus λ_0 and μ_0 are distinct eigenvalues of A . The eigenvectors $u(\lambda_0)$ and $u(\mu_0)$ are orthonormal:

$$u(\lambda_0)^T u(\mu_0) = f(\lambda_0, \mu_0) = 0.$$

Let λ and μ be approximate eigenvalues in the sense that

$$\begin{aligned} -\delta_j &= \frac{\lambda - \lambda_0}{\alpha_j - \lambda_0}, & |\delta_j| &\leq \frac{\delta}{1 + \delta}, \\ -\delta'_j &= \frac{\mu - \mu_0}{\alpha_j - \mu_0}, & |\delta'_j| &\leq \frac{\delta}{1 + \delta}. \end{aligned} \quad (12)$$

Here $\delta > 0$ is hopefully, but not necessarily, close to the machine unit ϵ . Note that (12) is equivalent with

$$\frac{\alpha_j - \lambda}{\alpha_j - \lambda_0} = 1 + \delta_j, \quad \frac{\alpha_j - \mu}{\alpha_j - \mu_0} = 1 + \delta'_j.$$

These conditions imply that the approximate eigenvectors $u(\lambda)$ and $u(\mu)$ are nearly orthogonal. For we have

$$\begin{aligned} \sqrt{f'(\lambda)f'(\mu)} u(\lambda)^T u(\mu) &= f(\lambda, \mu) - f(\lambda_0, \mu_0) \\ &= \sum_{j=1}^{n-1} \frac{\beta_j^2}{(\alpha_j - \lambda)(\alpha_j - \mu)} \left(1 - \frac{(\alpha_j - \lambda)(\alpha_j - \mu)}{(\alpha_j - \lambda_0)(\alpha_j - \mu_0)} \right) \\ &= \sum_{j=1}^{n-1} \frac{\beta_j^2}{(\alpha_j - \lambda)(\alpha_j - \mu)} (\delta_j + \delta'_j + \delta_j \delta'_j). \end{aligned}$$

Since

$$|\delta_j + \delta'_j + \delta_j \delta'_j| \leq \frac{2\delta}{1 + \delta} + \frac{\delta^2}{(1 + \delta)^2} \leq 2\delta$$

then

$$\sqrt{f'(\lambda)f'(\mu)} u(\lambda)^T u(\mu) = 2\delta \mathbf{b}^T (D - \lambda I)^{-1} \Theta (D - \mu I)^{-1} \mathbf{b}$$

with $|\Theta| \leq I$. Thus

$$\sqrt{f'(\lambda)f'(\mu)} |u(\lambda)^T u(\mu)| = 2\delta \| (D - \lambda I)^{-1} \mathbf{b} \|_2 \| (D - \mu I)^{-1} \mathbf{b} \|_2,$$

and so

$$|u(\lambda)^T u(\mu)| < 2\delta.$$

Condition (12) is stringent. If we let $\beta_k \rightarrow 0$ then it is easy to show that A can have an eigenvalue $\lambda_0 = \lambda_0(\beta_k) = \alpha_k + O(\beta_k^2)$; (12) then requires that the approximate eigenvalue λ satisfies a bound

$$|\lambda - \lambda_0| \leq O(\delta \beta_k^2),$$

which is difficult if $\beta_k/\|b\|$ is only somewhat larger than machine precision, say $\epsilon^{3/4}$. Two techniques are used to attempt to satisfy (12) - shifts of the origin [15], and simulated extended precision (SEP) arithmetic [26, 14]. Condition (12) means that

$$|\lambda - \lambda_0| < \delta \min\{\lambda_0 - \alpha_k, \alpha_{k-1} - \lambda_0\}.$$

When shifts are used it means that λ is nearly $f_l(\lambda_0)$.

4 Numerical stability of the algorithm

We now give a partial analysis of the stability of this approach to the eigenproblem for the symmetric arrow matrix. Observe that

$$f(\lambda) = \frac{p(\lambda)}{q(\lambda)} := \frac{\prod_{j=1}^n (\lambda - \lambda_j)}{\prod_{j=1}^{n-1} (\lambda - \alpha_j)}.$$

The following *inverse eigenvalue problem* [6] is important: given $\{\alpha_j\}$ and $\{\lambda_j\}$ satisfying (4), find $\{\beta_j\}$ and γ so that $\lambda(A) = \{\lambda_j\}$. This problem is simply solved by computing the residues of the partial fraction decomposition of f . In particular

$$\begin{aligned} \beta_k^2 &= \lim_{\lambda \rightarrow \alpha_k} (\alpha_k - \lambda) \frac{p(\lambda)}{q(\lambda)} \\ &= - \frac{\prod_{j=1}^n (\alpha_k - \lambda_j)}{\prod_{j \neq k} (\alpha_k - \alpha_j)}, \\ \gamma &= \sum_{j=1}^n \lambda_j - \sum_{j=1}^{n-1} \alpha_j. \end{aligned}$$

For fixed $\{\alpha_j\}$, the elements of the arrow *head*, $\{\beta_k\}$ and γ , are *explicitly known* functions of the eigenvalues.

Now let $\{\hat{\lambda}_j\}$ be a set of *approximate* eigenvalues of A satisfying (4). Then

$$\hat{\beta}_k^2 = - \frac{\prod_{j=1}^n (\alpha_k - \hat{\lambda}_j)}{\prod_{j \neq k} (\alpha_k - \alpha_j)} \quad (\hat{\beta}_k > 0), \quad (13)$$

$$\hat{\gamma} = \sum_{j=1}^n \hat{\lambda}_j - \sum_{j=1}^{n-1} \alpha_j. \quad (14)$$

define a *modified* matrix \hat{A} with $\lambda(\hat{A}) = \{\hat{\lambda}_j\}$. To obtain a backward error analysis for the complete eigenvalue problem we bound the differences $\hat{\beta}_k - \beta_k$ and $\hat{\gamma} - \gamma$.

4.1 Error analysis for the Dongarra-Sorensen condition

We give an error analysis using the Dongarra-Sorensen condition

$$-\frac{\tilde{\lambda}_j - \lambda_j}{\alpha_k - \lambda_j} = \delta_{j,k}, \quad |\delta_{j,k}| \leq \delta, \quad (15)$$

where $\delta = O(\epsilon)$ is of the order of the machine unit, simplifying that in [6].

Rearrangement of (15) gives

$$\tilde{\lambda}_j - \alpha_k = (\lambda_j - \alpha_k)(1 + \delta_{j,k}).$$

It follows that

$$\tilde{\beta}_k^2 = \beta_k^2 \prod_{j=1}^n (1 + \delta_{j,k}) = \beta_k^2 \left(1 + \sum_{j=1}^n \delta'_{j,k} \right),$$

and

$$\beta_k = \beta_k \left(1 + \frac{1}{2} \sum_{j=1}^n \delta''_{j,k} \right),$$

with the $\delta'_{j,k}$ and $\delta''_{j,k}$ at most only slightly larger than the $\delta_{j,k}$. Thus

$$\left| \frac{\beta_k - \beta_k}{\beta_k} \right| \leq \frac{n}{2} \delta'',$$

where $\delta'' = O(\epsilon)$ is only slightly larger than δ .

Now (14) becomes

$$\gamma = \gamma + \sum_{j=1}^n (\lambda_j - \alpha_{k(j)}) \delta_{j,k(j)}$$

with $\alpha_{k(j)}$ one of the poles of f . Thus

$$|\gamma - \gamma| \leq \delta \sum_{j=1}^n |\lambda_j - \alpha_{k(j)}|.$$

To minimize this bound we choose $\alpha_{k(j)}$ to be a pole of f closest to λ_j . Clearly, $\alpha_{k(1)} = \alpha_1$ and $\alpha_{k(n)} = \alpha_{n-1}$, so

$$|\tilde{\gamma} - \gamma| \leq \delta \left((\lambda_1 - \alpha_1) + \sum_{j=2}^{n-1} |\lambda_j - \alpha_{k(j)}| + (\alpha_{n-1} - \lambda_n) \right).$$

For $1 < j < n$ a closest pole to λ_j is either α_j or α_{j-1} . The distance

$$|\lambda_j - \alpha_{k(j)}| = \min \{ \lambda_j - \alpha_j, \alpha_{j-1} - \lambda_j \}$$

is maximized when λ_j is the midpoint of the interval (α_j, α_{j+1}) , and the value of the maximum is $(\alpha_j + \alpha_{j+1})/2$. Thus

$$\begin{aligned} |\hat{\gamma} - \gamma| &\leq \delta \left((\lambda_1 - \alpha_1) + \frac{1}{2} \sum_{j=2}^{n-1} (\alpha_{j+1} - \alpha_j) + (\alpha_{n+1} - \lambda_n) \right) \\ &= \delta \left((\lambda_1 - \lambda_n) - \frac{\alpha_1 - \alpha_{n+1}}{2} \right) \\ &\leq \delta(\lambda_1 - \lambda_n) \leq 2\delta \|A\|_2. \end{aligned}$$

In summary, the Dongarra-Sorensen condition implies small relative errors in each β_k and a small absolute error in γ . For the HSVD this implies small element-wise relative errors since the condition $\gamma = \gamma = 0$ is enforced by $\tilde{\lambda}_j + \tilde{\lambda}_{n+1-j} = 0$ (only half of the eigenvalues are actually computed, the rest follow from this condition).

4.2 Rounding error analysis of the computation of $f(\lambda)$

The choice of a termination criterion depends on a careful rounding error analysis of the particular manner in which we compute $f(\lambda)$. Let $\{\alpha_i\}$, $\{\beta_i\}$, and γ be floating point numbers. We represent λ as the ordered pair of floating point numbers (σ, μ) where the *shift* σ is a pole closest to λ , and $\lambda := \sigma + \mu$. For the exterior intervals we have $\sigma = \alpha_1$ or $\sigma = \alpha_{n+1}$. For the interior intervals σ can be determined by evaluating f at the midpoint and checking the sign. We compute $f(\lambda)$ as

$$f_\sigma(\mu) = \sum_{j=1}^{n-1} \frac{\beta_j^2}{\alpha'_j - \mu} + (\mu - \gamma').$$

with the standard operation precedence rules, where

$$\alpha'_j = \alpha_j - \sigma \quad \text{and} \quad \gamma' = \gamma - \sigma.$$

We use Wilkinson's notation: $fl(x * y) = (x * y)(1 + \delta)$ with $|\delta| \leq \epsilon/(1 + \epsilon)$ and $\epsilon = 2^{-t}$ the machine unit. More generally, ϵ denotes numbers not essentially larger than 2^{-t} [27] and the rounding errors δ satisfy $|\delta| < \epsilon$.

We define

$$fl(\alpha_j - \lambda) := fl(\alpha'_j - \mu) = fl((\alpha_j - \sigma) - \mu).$$

If $\sigma = \alpha_k$ then

$$fl(\alpha_k - \lambda) = -\mu = \alpha_k - \lambda,$$

with no rounding error. For $j \neq k$,

$$fl(\alpha_j - \lambda) = (\alpha_j - \lambda) \left(1 + \frac{\alpha_j - \alpha_k}{\alpha_j - \lambda} \delta + \delta' \right),$$

and since α_k is a pole closest to λ then $\left| \frac{\alpha_j - \alpha_k}{\alpha_j - \lambda} \right| \leq 2$. Thus all terms $\alpha_j - \lambda$ are computed with small relative errors:

$$fl(\alpha_j - \lambda) = (\alpha_j - \lambda)(1 + 3\delta_j), \quad |\delta_j| < \epsilon. \quad (16)$$

When computing $f(\lambda) = f_\sigma(\mu)$ we add the term $\lambda - \gamma = (\lambda - \sigma) - (\gamma - \sigma)$ last. A routine error analysis using (16) and

$$|\lambda - \gamma| \leq |f(\lambda)| + \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \lambda|}$$

to eliminate the term $|\lambda - \gamma|$ from the error bound gives

$$|fl(f(\lambda)) - f(\lambda)| \leq \epsilon \left(3|f(\lambda)| + |\sigma - \lambda| + (n+5) \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \lambda|} \right)$$

which implies

$$|fl(f(\lambda))| \leq (1+3\epsilon)|f(\lambda)| + \epsilon \left(|\sigma - \lambda| + (n+5) \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \lambda|} \right). \quad (17)$$

4.3 Termination

Our goal is to choose a termination criterion so that we stop when λ is as close to the true eigenvalue λ_k as possible. Let $\mu = \lambda_k$ in (11) with $f(\lambda_k) = 0$. Now $\alpha_k < \lambda_k < \alpha_{k+1}$. Also let $\alpha_k < \lambda < \alpha_{k+1}$. Then the terms $\alpha_j - \lambda$ and $\alpha_j - \lambda_k$ have the same sign and

$$|\lambda - \lambda_k| \leq \frac{|f(\lambda)|}{1 + \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \lambda||\alpha_j - \lambda_k|}}. \quad (18)$$

To obtain an upper bound for $|\lambda - \lambda_k|$ we need an upper bound for $|f(\lambda)|$ and a lower bound for the denominator. For the latter we have

$$1 + \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \lambda||\alpha_j - \lambda_k|} \geq 1 + \frac{\sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \lambda|}}{\max_j |\alpha_j - \lambda_k|}. \quad (19)$$

Let us determine how small $|f(\lambda)|$ is when λ is the rounded representation of λ_k . This is

$$\begin{aligned}\tilde{\lambda} &:= \sigma + f(\mu_k) = \sigma + \mu_k(1 + \delta) \\ &= \lambda_k + \mu_k\delta = \lambda_k + (\lambda_k - \sigma)\delta\end{aligned}$$

and we have

$$|\sigma - \lambda_k| = \min_j |\alpha_j - \lambda_k|.$$

Thus

$$\begin{aligned}|f(\tilde{\lambda})| &= |\tilde{\lambda} - \lambda_k| \left(1 + \sum_{j=1}^{n-1} \frac{\beta_j^2}{(\alpha_j - \tilde{\lambda})(\alpha_j - \lambda_k)} \right) \\ &= |(\sigma - \lambda_k)\delta| \left(1 + \sum_{j=1}^{n-1} \frac{\beta_j^2}{(\alpha_j - \tilde{\lambda})(\alpha_j - \lambda_k)} \right) \\ &\leq \epsilon \left(|\sigma - \lambda_k| + \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \tilde{\lambda}|} \right).\end{aligned}$$

From (17),

$$|f(f(\tilde{\lambda}))| \leq \epsilon \left(|\sigma - \lambda_k| + |\tilde{\lambda} - \sigma| + (n+6) \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \tilde{\lambda}|} \right).$$

Since $\lambda_k - \sigma = (\tilde{\lambda} - \sigma)/(1 + \delta)$ then

$$|f(f(\tilde{\lambda}))| \leq \epsilon \left(2|\tilde{\lambda} - \sigma| + (n+6) \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \tilde{\lambda}|} \right).$$

We terminate and set $\lambda_k := \lambda$ when

$$|f(f(\lambda))| \leq 2\epsilon \left(2|\lambda - \sigma| + (n+6) \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \lambda|} \right).$$

Inequality (17) also holds if $f(\lambda)$ and $f(f(\lambda))$ are interchanged. Thus

$$|f(\tilde{\lambda}_k)| \leq \epsilon \left(5|\tilde{\lambda}_k - \sigma| + (3n+17) \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \tilde{\lambda}_k|} \right). \quad (20)$$

From (18) and (19)

$$|\tilde{\lambda}_k - \lambda_k| \leq \epsilon \max_j |\lambda_k - \alpha_j| \frac{5|\sigma - \tilde{\lambda}_k| + (3n+17) \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \tilde{\lambda}_k|}}{\max_j |\lambda_k - \alpha_j| + \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \lambda_k|}}.$$

Since $|\sigma - \tilde{\lambda}_k| \leq |\sigma - \lambda_k| + |\tilde{\lambda}_k - \lambda_k|$ and $|\sigma - \lambda_k| \leq \max_j |\alpha_j - \lambda_k|$ the computed eigenvalues satisfy

$$\begin{aligned} |\lambda_k - \tilde{\lambda}_k| &\leq (3n+17)\epsilon \max_j |\alpha_j - \lambda_k| \\ &\leq 6(n+6)\epsilon \|A\|_2. \end{aligned} \quad (21)$$

4.4 Error analysis for the Gu-Eisenstat condition

From $\tilde{\gamma} - \gamma = \sum_{j=1}^n (\lambda_j - \tilde{\lambda}_j)$ and (21) we find

$$|\tilde{\gamma} - \gamma| \leq 6n(n+6)\epsilon \|A\|_2.$$

We have noted that the Dongarra-Sorensen condition (15) is stringent. It is natural to ask for small *absolute* errors in the β_k . If we replace $\delta_{j,k}$ by $\delta_{j,k}/\beta_k$ in the analysis in section 4.1 we find that

$$\beta_k = \beta_k \left(1 + \frac{1}{2} \sum_{j=1}^{n-1} \frac{\delta_{j,k}''}{\beta_k} \right) = \beta_k + \frac{1}{2} \sum_{j=1}^{n-1} \delta_{j,k}'',$$

and

$$|\beta_k - \tilde{\beta}_k| \leq \frac{n}{2} \epsilon'', \quad |\delta''| \leq \epsilon(1 + O(\epsilon)),$$

are implied by the *Gu-Eisenstat condition*

$$-\beta_k \frac{\lambda_j - \tilde{\lambda}_j}{\alpha_j - \tilde{\lambda}_j} = \delta_{j,k}, \quad |\delta_{j,k}| \leq \delta.$$

We must bound δ .

From (20)

$$|\tilde{\lambda}_k - \lambda_k| \left(1 + \sum_{j=1}^{n-1} \frac{\beta_j^2}{(\alpha_j - \lambda_k)(\alpha_j - \tilde{\lambda}_k)} \right) \leq m \left(|\lambda_k - \sigma| + \sum_{j=1}^{n-1} \frac{\beta_j^2}{|\alpha_j - \tilde{\lambda}_k|} \right)$$

with $m = 3(n+6)$. Using

$$|\lambda - \sigma| \leq |\lambda - \lambda_k| + |\lambda_k - \sigma|$$

and the Gu-Eisenstat inequality,

$$\frac{1}{|\alpha_j - \lambda|} \leq \left| \frac{1}{(\alpha_j - \lambda)(\alpha_j - \lambda_k)} \right|^{1/2} + \frac{|\lambda_k - \lambda|}{(\alpha_j - \lambda)(\alpha_j - \lambda_k)},$$

we get

$$|\hat{\lambda}_k - \lambda_k| \leq m\epsilon \left(|\lambda_k - \sigma| + \frac{\sum_{j=1}^{n-1} \frac{\beta_j^2}{[(\alpha_j - \hat{\lambda}_k)(\alpha_j - \lambda_k)]^{1/2}}}{\sum_{j=1}^{n-1} \frac{\beta_j^2}{(\alpha_j - \lambda_k)(\alpha_j - \lambda_k)}} \right),$$

where ϵ has been increased to $\epsilon/(1 - m\epsilon)$.

By Cauchy's inequality,

$$\begin{aligned} |\hat{\lambda}_k - \lambda_k| &\leq m\epsilon \left(|\lambda_k - \sigma| + \frac{\|\mathbf{b}\|}{\left(\sum_{j=1}^{n-1} \frac{\beta_j^2}{(\alpha_j - \lambda_k)(\alpha_j - \lambda_k)}\right)^{1/2}} \right) \\ &\leq m\epsilon \left(|\lambda_k - \sigma| + \frac{\|\mathbf{b}\|}{\beta_j} \left[(\alpha_j - \hat{\lambda}_k)(\alpha_j - \lambda_k)\right]^{1/2} \right), \end{aligned}$$

for every j . The arithmetic-geometric mean inequality and the triangle inequality yield

$$|\hat{\lambda}_k - \lambda_k| \leq m\epsilon \left(|\lambda_k - \sigma| + \frac{\|\mathbf{b}\|}{\beta_j} \left(|\alpha_j - \lambda_k| + \frac{1}{2} |\lambda_k - \hat{\lambda}_k| \right) \right).$$

Thus

$$\begin{aligned} \left(1 - \frac{m\epsilon \|\mathbf{b}\|}{\beta_j}\right) |\hat{\lambda}_k - \lambda_k| &\leq m\epsilon \left(|\lambda_k - \sigma| + |\alpha_j - \lambda_k| \frac{\|\mathbf{b}\|}{\beta_j} \right) \\ &= m\epsilon \|\mathbf{b}\| \frac{|\alpha_j - \lambda_k|}{\beta_j} \left(1 + \frac{|\lambda_k - \sigma|}{|\lambda_k - \alpha_j|} \frac{\beta_j}{\|\mathbf{b}\|} \right) \\ &\leq 2m\epsilon \|\mathbf{b}\| \frac{|\alpha_j - \lambda_k|}{\beta_j}. \end{aligned}$$

If $m\epsilon \|\mathbf{b}\| \leq \beta_j$ for all j , then

$$|\delta_{j,k}| \leq 4m\epsilon \|\mathbf{b}\|.$$

and consequently

$$|\beta_k - \beta_k| \leq 6n(n+6)\epsilon \|\mathbf{b}\|.$$

Thus tol_β is $m\epsilon$. If $\beta_k < 3(n+6)\epsilon \|\mathbf{b}\|$ we replace β_k by zero and accept α_k as an eigenvalue with normalized eigenvector \mathbf{e}_k .

The computed eigenvectors of A are taken to be those of the nearby matrix \hat{A} . Because of (13) and (16) they are computed to high relative precision elementwise and hence are numerically orthogonal [20].

5 Acknowledgements

We wish to thank Ming Gu and Stan Eisenstat for providing a preprint of their manuscript [20]. It is our understanding that they have independently extended their results to include the arrow case in [21]. Both authors were supported by direct grant from the Naval Postgraduate School. The second author also acknowledges support from the Interdisciplinary Project Center for Supercomputing at the ETH, Zürich.

References

- [1] N. I. AKHIEZER, *The classical moment problem and some related questions in analysis*, Hafner Publishing Company, 1961.
- [2] G. S. AMMAR, L. REICHEL, AND D. C. SORENSEN, *An implementation of a divide and conquer algorithm for the unitary eigenproblem*, ACM Trans. Math. Software, 18 (1992), pp. 292-307.
- [3] P. ARBENZ, *Computing eigenvalues of banded symmetric Toeplitz matrices*, SIAM J. Sci. Statist. Comput., 12 (1991), pp. 743-754.
- [4] ———, *Divide and conquer algorithms for the bandsymmetric eigenvalue problem*, in Parallel Computing '91, D. I. Evans, G. R. Joubert, and H. Liddell, eds., Elsevier Science Publishers B. V., Amsterdam, 1992, pp. 151-158.
- [5] P. ARBENZ AND G. H. GOLUB, *On the spectral decomposition of hermitian matrices modified by low rank perturbations with applications*, SIAM J. Matrix Anal. Appl., 9 (1988), pp. 40-58.
- [6] J. L. BARLOW, *Error analysis of update methods for the symmetric eigenvalue problem*, SIAM J. Matrix Anal. Appl. To appear.
- [7] C. BEATTIE AND D. W. LON, *Schur complements and the Weinstein-Aronszajn theory for modified matrix eigenvalue problems*, Linear Algebra Appl., 108 (1988), pp. 37-61.
- [8] C. BEATTIE AND C. J. HIBBENS, *Parallel solution of a generalized symmetric matrix eigenvalue problem*, Working paper, Department of Mathematics, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, 1992.
- [9] A. A. BEEX AND M. P. FARGUES, *Highly parallel recursive/iterative Toeplitz eigenspace decomposition*, IEEE Trans. Acoustics, Speech, Signal Proc., 37 (1989), pp. 1765-1768.
- [10] A. A. BEEX, D. M. WILKES, AND M. P. FARGUES, *The C-RISE algorithm and the generalized eigenvalue problem*, Signal Processing, Submitted.
- [11] J. R. BUNCH AND C. NIELSEN, *Updating the singular value decomposition*, Numer. Math., 31 (1978), pp. 111-129.

- [12] J. R. BUNCH, C. NIELSEN, AND D. C. SORENSEN, *Rank one modification of the symmetric eigenproblem*, Numer. Math., 31 (1978), pp. 31-48.
- [13] J. J. M. CUPPEN, *A divide and conquer method for the symmetric tridiagonal eigenproblem*, Numer. Math., 36 (1981), pp. 177-195.
- [14] T. J. DEKKER, *A floating-point technique for extending the available precision*, Numer. Math., 18 (1971), pp. 224-242.
- [15] J. J. DONGARRA AND D. C. SORENSEN, *A fully parallel algorithm for the symmetric eigenvalue problem*, SIAM J. Sci. Statist. Comput., 8 (1987), pp. 139-154.
- [16] G. GOLUB AND W. KAHAN, *Calculating the singular values and pseudo-inverse of a matrix*, J. Soc. Indust. Appl. Math. Ser. B Numer. Anal., 2 (1965).
- [17] G. H. GOLUB, *Some modified matrix eigenvalue problems*, SIAM Rev., 15 (1973), pp. 318-334.
- [18] W. B. GRAGG AND L. REICHEL, *A divide and conquer method for unitary and orthogonal eigenproblems*, Numer. Math., 57 (1990), pp. 695-718.
- [19] W. B. GRAGG, J. R. THORNTON, AND D. D. WARNER, *Parallel divide and conquer algorithms for the symmetric tridiagonal eigenproblem and bidiagonal singular value problem*, in Modeling and Simulation, vol. 23, part 1, W. G. Vogt and M. H. Mickle, eds., Univ. Pittsburgh School of Engineering, 1992, pp. 49-56.
- [20] M. GU AND S. C. EISENSTAT, *A stable and efficient algorithm for the rank-one modification of the symmetric eigenproblem*, SIAM J. Matrix Anal. Appl. To appear.
- [21] M. GU AND S. C. EISENSTAT, *A divide-and-conquer algorithm for the symmetric tridiagonal eigenproblem*, Working paper, Department of Computer Science, Yale University (1992).
- [22] E. R. JESSUP AND D. C. SORENSEN, *A parallel algorithm for computing the singular value decomposition of a matrix*, SIAM J. Matrix Anal. Appl. To appear.
- [23] R.-C. LI, *Solving secular equations stably and efficiently*, Working paper, Department of Mathematics, University of California, Berkeley, (1992).
- [24] K. LÖWNER, *Über monotone Matrixfunktionen*, Math. Z., 38 (1934), pp. 177-216.
- [25] D. P. O'LEARY AND G. W. STEWART, *Computing the eigenvalues and eigenvectors of arrowhead matrices*, J. Comp. Phys., 90 (1990), pp. 497-505.
- [26] D. C. SORENSEN AND P. T. TANG, *On the orthogonality of eigenvectors computed by divide and conquer techniques*, SIAM J. Numer. Anal., 28 (1991), pp. 1752-1775.
- [27] J. H. WILKINSON, *Rounding errors in algebraic processes*, Prentice-Hall, 1963.
- [28] J. H. WILKINSON, *The algebraic eigenvalue problem*, Oxford University Press, 1965.

DISTRIBUTION LIST

Director (2)
Defense Tech Information Center
Cameron Station
Alexandria, VA 22314

Research Office (1)
Code 81
Naval Postgraduate School
Monterey, CA 93943

Library (2)
Code 52
Naval Postgraduate School
Monterey, CA 93943

Professor Richard Franke (1)
Department of Mathematics
Naval Postgraduate School
Monterey, CA 93943

Dr. Neil L. Gerr (1)
Mathematical Sciences Division
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217-5000

Dr. Richard Lau (1)
Mathematical Sciences Division
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217-5000

Harper Whitehouse (Code 743) (1)
NCCOSC RDT&E Division
271 Catalina Blvd.
San Diego, CA 92152-5000

Keith Bromley (Code 7601) (1)
NCCOSC RDT&E Division
271 Catalina Blvd.
San Diego, CA 92152-5000

John Rockway (Code 804) (1)
NCCOSC RDT&E Division
271 Catalina Blvd.
San Diego, CA 92152-5000

Professor Carlos Borges (15)
Department of Mathematics
Naval Postgraduate School
Monterey, CA 93943